

ADVANCED GCE PHYSICS A

G484

The Newtonian World

Candidates answer on the Question Paper

OCR Supplied Materials:

Data, Formulae and Relationships Booklet

Other Materials Required:

Electronic calculator

Tuesday 29 June 2010 Afternoon

Duration: 1 hour



Candidate Forename				Candidate Surname			
Centre Number	r			Candidate N	umber		

INSTRUCTIONS TO CANDIDATES


- Write your name clearly in capital letters, your Centre Number and Candidate Number in the boxes above.
- Use black ink. Pencil may be used for graphs and diagrams only.
- Read each question carefully and make sure that you know what you have to do before starting your answer.
- Answer all the questions.
- Do not write in the bar codes.
- Write your answer to each question in the space provided. Additional paper may be used if necessary but you must clearly show your Candidate Number, Centre Number and question number(s).

INFORMATION FOR CANDIDATES

- The number of marks is given in brackets [] at the end of each question or part question.
- The total number of marks for this paper is 60.
- You may use an electronic calculator.
- You are advised to show all the steps in any calculations.
- Where you see this icon you will be awarded marks for the quality of written communication in your answer.

This means for example you should:

- ensure that text is legible and that spelling, punctuation and grammar are accurate so that meaning is
- organise information clearly and coherently, using specialist vocabulary when appropriate.
- This document consists of 12 pages. Any blank pages are indicated.



Answer all the questions.

(a) A particular collision between two objects is *inelastic*. Place a tick (✓) at the end of each statement that applies to such a collision.

Statement	
The magnitude of the impulse on each object is the same.	
Kinetic energy and momentum for the objects are conserved.	
Total energy is conserved.	
After the collision, the objects have the same momentum.	

(b) Fig. 1.1 shows a tennis ball before and after striking a wall at right angles.




Fig. 1.1

The ball of mass 0.060 kg hits the wall at a speed of 12 m s⁻¹. The ball is in contact with the wall for 0.15 s. It rebounds with a speed of 9.0 m s⁻¹. Calculate

(i) the loss of kinetic energy during the collision

(ii) the magnitude of the average force exerted on the ball by the wall

(iii) the magnitude of the average force exerted on the wall by the ball during this collision.

		average force on wall = N [1]
(c)	(i)	State three assumptions of the kinetic model of ideal gases.
		[3]
	(ii)	Use the kinetic theory of gases to explain how a gas exerts a pressure.
		[3] [Total: 13]
		[Total: 10]

2 (a) Fig. 2.1 shows an aeroplane flying in a horizontal circle at constant speed. The weight of the aeroplane is W and L is the lift force acting at right angles to the wings.




Fig. 2.1

(i)	Explain how the lift force L maintains the aeroplane flying in a horizontal circle.
	[2]
(ii)	The aeroplane of mass $1.2 \times 10^5 \mathrm{kg}$ is flying in a horizontal circle of radius 2.0 km.
	The centripetal force acting on the aeroplane is $1.8\times10^6\mathrm{N}$. Calculate the speed of the aeroplane.

speed = ms⁻¹ [2]

(b) Fig. 2.2 shows a satellite orbiting the Earth at a constant speed *v*. The radius of the orbit is *r*.




Fig. 2.2

Show that the orbital period T of the satellite is given by the equation

$$T^2 = \frac{4\pi^2 r^3}{GM}$$

where M is the mass of the Earth and G is the gravitational constant.

	ГЭ	•
	1.5	
U	L	

(c) The satellites used in television communication systems are usually placed in geostationary orbits.



In your answer, you should use appropriate technical words spelled correctly.

(i)	State two features of geostationary orbits.
	1
	2
	[2]
	<u></u>

(ii) Calculate the radius of orbit of a geostationary satellite.

The mass of the Earth is 6.0×10^{24} kg.

[Total: 12]

3	(a)	State two conditions concerning the acceleration of an oscillating object that must apply for
		simple harmonic motion.

1.	
• • •	
2.	
• •	
	[2]

(b) Fig. 3.1 shows how the potential energy, in mJ, of a simple harmonic oscillator varies with displacement.




Fig. 3.1

On Fig. 3.1 sketch graphs to show the variation of

- (i) kinetic energy of the oscillator with displacement label this graph K
- (ii) the total energy of the oscillator with displacement label this graph T. [1]

[2]

(c)	Use	e Fig. 3.1 to determine
	(i)	the amplitude of the oscillations
		amplitude = m [1]
	(ii)	the maximum speed of the oscillator of mass 0.12 kg
		maximum speed = ms ⁻¹ [2]
	(iii)	the frequency of the oscillations.
		frequency = Hz [2]
(-1)	D	
(u)	a u	sonance can either be useful or a problem. Describe one example where resonance has seful application and one example where resonance is a problem or nuisance. For each mple identify what is oscillating and what causes these oscillations.
	(i)	useful application
		[2]
	(ii)	problem
		[2]

[Total: 14]

4 Fig. 4.1 shows smoke particles suspended in air. The arrows indicate the directions in which the smoke particles are moving at a particular instant. The lengths of the arrows indicate the different speeds of the particles.




Fig. 4.1

(a) (i)	State the name given to this type of random motion of smoke particles in air.
	In your answer, you should use appropriate technical terms spelled correctly.
	[1]
(ii)	State two conclusions about the air molecules that may be deduced from the observed motion of the smoke particles.
	[2]

(b)	(i)		radius of an inflated football is 0.11 m. The temperature of the air inside the ball is C. Calculate the mass of air in the ball when the pressure inside it is $2.6 \times 10^5 \text{Pa}$.
		The	mass of one mole of air is 0.028 kg.
			mass of air = kg [4]
	(ii)	The	football is left in a room at a temperature of 0 °C until it reaches thermal equilibrium.
		1	Explain the term thermal equilibrium.
			[1]
		2	Calculate the pressure exerted by the air inside the football when the temperature drops to 0 $^{\circ}\text{C}.$
			pressure = Pa [2]
			[Total: 10]

© OCR 2010 Turn over

		mass $970\mathrm{kg}$ is travelling at $27\mathrm{ms^{-1}}$ when the brakes are applied. The car is brought to distance of $40\mathrm{m}$.
(a)	(i)	Calculate the kinetic energy of the car when it is travelling at 27 m s ⁻¹ .
		kinetic energy = J [1]
	(ii)	Hence calculate the average braking force on the car stating any assumption that you make.
		average braking force = N
		assumption
(b)		
(b)		assumption
(b)	ma	assumption

(ii)	State and explain two reasons why the actual temperature rise will be different.
	[4]
(iii)	Suggest one modification to the design of the disc braking system that could reduce the temperature rise of the discs.
	[1]
	[Total: 11]

END OF QUESTION PAPER

PLEASE DO NOT WRITE ON THIS PAGE



Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations, is given to all schools that receive assessment material and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

 $For queries \ or \ further \ information \ please \ contact \ the \ Copyright \ Team, \ First \ Floor, 9 \ Hills \ Road, \ Cambridge \ CB2 \ 1GE.$

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

© OCR 2010