



**Sixth Term Examination Papers**  
**MATHEMATICS 3**  
**Friday 21 June 2019**

**9475**  
Morning  
Time: 3 hours

Additional Material: Answer Booklet

**INSTRUCTIONS TO CANDIDATES**

Read this page carefully, but do not open this question paper until you are told that you may do so.

Read the additional instructions on the front of the answer booklet.

Write your name, centre number, candidate number, date of birth, and indicate the paper number in the spaces provided on the answer booklet.

Make sure you fill in page 1 **AND** page 3 of the answer booklet with your details.

**INFORMATION FOR CANDIDATES**

There are 12 questions in this paper.

Each question is marked out of 20. There is no restriction of choice.

All questions attempted will be marked.

Your final mark will be based on the **six** questions for which you gain the highest marks.

You are advised to concentrate on no more than **six** questions. Little credit will be given for fragmentary answers.

**There is NO Mathematical Formulae booklet.**

**Calculators are not permitted.**

**Wait to be told you may begin before turning this page.**

---

This question paper consists of 9 printed pages and 3 blank pages.



## ERRATUM

### Question 5 part (ii)

The sentence

“Use the substitution  $y = \frac{bx}{\sqrt{x^2+p}}$ , where p is a suitably chosen constant, ...”

Should read

“Use the substitution  $y = \frac{cx}{\sqrt{x^2+p}}$ , where p is a suitably chosen constant, ...”

## Section A: Pure Mathematics

- 1 The coordinates of a particle at time  $t$  are  $x$  and  $y$ . For  $t \geq 0$ , they satisfy the pair of coupled differential equations

$$\begin{aligned}\dot{x} &= -x - ky \\ \dot{y} &= x - y\end{aligned}$$

where  $k$  is a constant. When  $t = 0$ ,  $x = 1$  and  $y = 0$ .

- (i) Let  $k = 1$ . Find  $x$  and  $y$  in terms of  $t$  and sketch  $y$  as a function of  $t$ .

Sketch the path of the particle in the  $x$ - $y$  plane, giving the coordinates of the point at which  $y$  is greatest and the coordinates of the point at which  $x$  is least.

- (ii) Instead, let  $k = 0$ . Find  $x$  and  $y$  in terms of  $t$  and sketch the path of the particle in the  $x$ - $y$  plane.

- 2 The definition of the derivative  $f'$  of a (differentiable) function  $f$  is

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}. \quad (*)$$

- (i) The function  $f$  has derivative  $f'$  and satisfies

$$f(x+y) = f(x)f(y)$$

for all  $x$  and  $y$ , and  $f'(0) = k$  where  $k \neq 0$ . Show that  $f(0) = 1$ .

Using  $(*)$ , show that  $f'(x) = kf(x)$  and find  $f(x)$  in terms of  $x$  and  $k$ .

- (ii) The function  $g$  has derivative  $g'$  and satisfies

$$g(x+y) = \frac{g(x) + g(y)}{1 + g(x)g(y)}$$

for all  $x$  and  $y$ ,  $|g(x)| < 1$  for all  $x$ , and  $g'(0) = k$  where  $k \neq 0$ .

Find  $g'(x)$  in terms of  $g(x)$  and  $k$ , and hence find  $g(x)$  in terms of  $x$  and  $k$ .

3 The matrix  $\mathbf{A}$  is given by

$$\mathbf{A} = \begin{pmatrix} a & b \\ c & d \end{pmatrix}.$$

- (i) You are given that the transformation represented by  $\mathbf{A}$  has a line  $L_1$  of invariant points (so that each point on  $L_1$  is transformed to itself). Let  $(x, y)$  be a point on  $L_1$ . Show that  $((a - 1)(d - 1) - bc)xy = 0$ .

Show further that  $(a - 1)(d - 1) = bc$ .

What can be said about  $\mathbf{A}$  if  $L_1$  does not pass through the origin?

- (ii) By considering the cases  $b \neq 0$  and  $b = 0$  separately, show that if  $(a - 1)(d - 1) = bc$  then the transformation represented by  $\mathbf{A}$  has a line of invariant points. You should identify the line in the different cases that arise.
- (iii) You are given instead that the transformation represented by  $\mathbf{A}$  has an invariant line  $L_2$  (so that each point on  $L_2$  is transformed to a point on  $L_2$ ) and that  $L_2$  does not pass through the origin. If  $L_2$  has the form  $y = mx + k$ , show that  $(a - 1)(d - 1) = bc$ .

4 The  $n$ th degree polynomial  $P(x)$  is said to be *reflexive* if:

- (a)  $P(x)$  is of the form  $x^n - a_1x^{n-1} + a_2x^{n-2} - \dots + (-1)^na_n$  where  $n \geq 1$ ;  
(b)  $a_1, a_2, \dots, a_n$  are real;  
(c) the  $n$  (not necessarily distinct) roots of the equation  $P(x) = 0$  are  $a_1, a_2, \dots, a_n$ .
- (i) Find all reflexive polynomials of degree less than or equal to 3.
- (ii) For a reflexive polynomial with  $n > 3$ , show that

$$2a_2 = -a_2^2 - a_3^2 - \dots - a_n^2.$$

Deduce that, if all the coefficients of a reflexive polynomial of degree  $n$  are integers and  $a_n \neq 0$ , then  $n \leq 3$ .

- (iii) Determine all reflexive polynomials with integer coefficients.

5 (i) Let

$$f(x) = \frac{x}{\sqrt{x^2 + p}},$$

where  $p$  is a non-zero constant. Sketch the curve  $y = f(x)$  for  $x \geq 0$  in the case  $p > 0$ .

(ii) Let

$$I = \int \frac{1}{(b^2 - y^2)\sqrt{c^2 - y^2}} dy,$$

where  $b$  and  $c$  are positive constants. Use the substitution  $y = \frac{bx}{\sqrt{x^2 + p}}$ , where  $p$  is a suitably chosen constant, to show that

$$I = \int \frac{1}{b^2 + (b^2 - c^2)x^2} dx.$$

Evaluate

$$\int_1^{\sqrt{2}} \frac{1}{(3 - y^2)\sqrt{2 - y^2}} dy.$$

[ **Note:**  $\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a} + \text{constant.}$  ]

Hence evaluate

$$\int_{\frac{1}{\sqrt{2}}}^1 \frac{y}{(3y^2 - 1)\sqrt{2y^2 - 1}} dy.$$

(iii) By means of a suitable substitution, evaluate

$$\int_{\frac{1}{\sqrt{2}}}^1 \frac{1}{(3y^2 - 1)\sqrt{2y^2 - 1}} dy.$$

- 6** The point  $P$  in the Argand diagram is represented by the complex number  $z$ , which satisfies

$$zz^* - az^* - a^*z + aa^* - r^2 = 0.$$

Here,  $r$  is a positive real number and  $r^2 \neq a^*a$ . By writing  $|z - a|^2$  as  $(z - a)(z - a)^*$ , show that the locus of  $P$  is a circle,  $C$ , the radius and the centre of which you should give.

- (i) The point  $Q$  is represented by  $\omega$ , and is related to  $P$  by  $\omega = \frac{1}{z}$ . Let  $C'$  be the locus of  $Q$ . Show that  $C'$  is also a circle, and give its radius and centre.

If  $C$  and  $C'$  are the same circle, show that

$$(|a|^2 - r^2)^2 = 1$$

and that either  $a$  is real or  $a$  is imaginary. Give sketches to indicate the position of  $C$  in these two cases.

- (ii) Suppose instead that the point  $Q$  is represented by  $\omega$ , where  $\omega = \frac{1}{z^*}$ . If the locus of  $Q$  is  $C$ , is it the case that either  $a$  is real or  $a$  is imaginary?

- 7** The *Devil's Curve* is given by

$$y^2(y^2 - b^2) = x^2(x^2 - a^2),$$

where  $a$  and  $b$  are positive constants.

- (i) In the case  $a = b$ , sketch the Devil's Curve.
- (ii) Now consider the case  $a = 2$  and  $b = \sqrt{5}$ , and  $x \geq 0, y \geq 0$ .
- (a) Show by considering a quadratic equation in  $x^2$  that either  $0 \leq y \leq 1$  or  $y \geq 2$ .
- (b) Describe the curve very close to and very far from the origin.
- (c) Find the points at which the tangent to the curve is parallel to the  $x$ -axis and the point at which the tangent to the curve is parallel to the  $y$ -axis.

Sketch the Devil's Curve in this case.

- (iii) Sketch the Devil's Curve in the case  $a = 2$  and  $b = \sqrt{5}$  again, but with  $-\infty < x < \infty$  and  $-\infty < y < \infty$ .

- 8 A pyramid has a horizontal rectangular base  $ABCD$  and its vertex  $V$  is vertically above the centre of the base. The acute angle between the face  $AVB$  and the base is  $\alpha$ , the acute angle between the face  $BVC$  and the base is  $\beta$  and the obtuse angle between the faces  $AVB$  and  $BVC$  is  $\pi - \theta$ .

- (i) The edges  $AB$  and  $BC$  are parallel to the unit vectors  $\mathbf{i}$  and  $\mathbf{j}$ , respectively, and the unit vector  $\mathbf{k}$  is vertical. Find a unit vector that is perpendicular to the face  $AVB$ .

Show that

$$\cos \theta = \cos \alpha \cos \beta.$$

- (ii) The edge  $BV$  makes an angle  $\phi$  with the base. Show that

$$\cot^2 \phi = \cot^2 \alpha + \cot^2 \beta.$$

Show also that

$$\cos^2 \phi = \frac{\cos^2 \alpha + \cos^2 \beta - 2 \cos^2 \theta}{1 - \cos^2 \theta} \geq \frac{2 \cos \theta - 2 \cos^2 \theta}{1 - \cos^2 \theta}$$

and deduce that  $\phi < \theta$ .

## Section B: Mechanics

- 9 In this question,  $\mathbf{i}$  and  $\mathbf{j}$  are perpendicular unit vectors and  $\mathbf{j}$  is vertically upwards.

A smooth hemisphere of mass  $M$  and radius  $a$  rests on a smooth horizontal table with its plane face in contact with the table. The point  $A$  is at the top of the hemisphere and the point  $O$  is at the centre of its plane face.

Initially, a particle  $P$  of mass  $m$  rests at  $A$ . It is then given a small displacement in the positive  $\mathbf{i}$  direction. At a later time  $t$ , when the particle is still in contact with the hemisphere, the hemisphere has been displaced by  $-s\mathbf{i}$  and  $\angle AOP = \theta$ .

- (i) Let  $\mathbf{r}$  be the position vector of the particle at time  $t$  with respect to the initial position of  $O$ . Write down an expression for  $\mathbf{r}$  in terms of  $a$ ,  $\theta$  and  $s$  and show that

$$\dot{\mathbf{r}} = (a\dot{\theta} \cos \theta - \dot{s})\mathbf{i} - a\dot{\theta} \sin \theta \mathbf{j}.$$

Show also that

$$\dot{s} = (1 - k)a\dot{\theta} \cos \theta,$$

where  $k = \frac{M}{m + M}$ , and deduce that

$$\dot{\mathbf{r}} = a\dot{\theta}(k \cos \theta \mathbf{i} - \sin \theta \mathbf{j}).$$

- (ii) Show that

$$a\dot{\theta}^2 (k \cos^2 \theta + \sin^2 \theta) = 2g(1 - \cos \theta).$$

- (iii) At time  $T$ , when  $\theta = \alpha$ , the particle leaves the hemisphere. By considering the component of  $\ddot{\mathbf{r}}$  parallel to the vector  $\sin \theta \mathbf{i} + k \cos \theta \mathbf{j}$ , or otherwise, show that at time  $T$

$$a\dot{\theta}^2 = g \cos \alpha.$$

Find a cubic equation for  $\cos \alpha$  and deduce that  $\cos \alpha > \frac{2}{3}$ .



- 10** Two identical smooth spheres  $P$  and  $Q$  can move on a smooth horizontal table. Initially,  $P$  moves with speed  $u$  and  $Q$  is at rest. Then  $P$  collides with  $Q$ . The direction of travel of  $P$  before the collision makes an acute angle  $\alpha$  with the line joining the centres of  $P$  and  $Q$  at the moment of the collision. The coefficient of restitution between  $P$  and  $Q$  is  $e$  where  $e < 1$ . As a result of the collision,  $P$  has speed  $v$  and  $Q$  has speed  $w$ , and  $P$  is deflected through an angle  $\theta$ .

- (i) Show that

$$u \sin \alpha = v \sin(\alpha + \theta)$$

and find an expression for  $w$  in terms of  $v$ ,  $\theta$  and  $\alpha$ .

- (ii) Show further that

$$\sin \theta = \cos(\theta + \alpha) \sin \alpha + e \sin(\theta + \alpha) \cos \alpha$$

and find an expression for  $\tan \theta$  in terms of  $\tan \alpha$  and  $e$ .

Find, in terms of  $e$ , the maximum value of  $\tan \theta$  as  $\alpha$  varies.

## Section C: Probability and Statistics

- 11** The number of customers arriving at a builders' merchants each day follows a Poisson distribution with mean  $\lambda$ . Each customer is offered some free sand. The probability of any given customer taking the free sand is  $p$ .

- (i) Show that the number of customers each day who take sand follows a Poisson distribution with mean  $p\lambda$ .
- (ii) The merchant has a mass  $S$  of sand at the beginning of the day. Each customer who takes the free sand gets a proportion  $k$  of the remaining sand, where  $0 \leq k < 1$ . Show that by the end of the day the expected mass of sand taken is

$$(1 - e^{-kp\lambda})S.$$

- (iii) At the beginning of the day, the merchant's bag of sand contains a large number of grains, exactly one of which is made from solid gold. At the end of the day, the merchant's assistant takes a proportion  $k$  of the remaining sand. Find the probability that the assistant takes the golden grain. Comment on the case  $k = 0$  and on the limit  $k \rightarrow 1$ .

In the case  $p\lambda > 1$  find the value of  $k$  which maximises the probability that the assistant takes the golden grain.

- 12** The set  $S$  is the set of all integers from 1 to  $n$ . The set  $T$  is the set of all distinct subsets of  $S$ , including the empty set  $\emptyset$  and  $S$  itself. Show that  $T$  contains exactly  $2^n$  sets.

The sets  $A_1, A_2, \dots, A_m$ , which are not necessarily distinct, are chosen randomly and independently from  $T$ , and for each  $k$  ( $1 \leq k \leq m$ ), the set  $A_k$  is equally likely to be any of the sets in  $T$ .

- (i) Write down the value of  $P(1 \in A_1)$ .
- (ii) By considering each integer separately, show that  $P(A_1 \cap A_2 = \emptyset) = \left(\frac{3}{4}\right)^n$ .  
Find  $P(A_1 \cap A_2 \cap A_3 = \emptyset)$  and  $P(A_1 \cap A_2 \cap \dots \cap A_m = \emptyset)$ .
- (iii) Find  $P(A_1 \subseteq A_2)$ ,  $P(A_1 \subseteq A_2 \subseteq A_3)$  and  $P(A_1 \subseteq A_2 \subseteq \dots \subseteq A_m)$ .

**BLANK PAGE**

**BLANK PAGE**